

For guidance on planning your revegetation or restoration site (size, shape, species, density of planting) refer to chapters 4 and 5 of this guide, and head to www.revegetation.org.au

Remember: good quality vegetation sites may not need revegetation at all. Appropriate management can encourage natural regeneration.

TEN MILE



LANDFORM	Flats	Lower slopes	Hills – N and NW aspect		Hills – S and SE aspect
VEGETATION TYPE	Blakely's Red Gum woodland	Blakely's Red Gum and Yellow Box woodland	Dry forest		Moist open forest
GEOLOGY & SOILS	Alluvium – sand, silt, clay and gravel. Sandy yellow earths.	Alluvium – sand, silt, clay and gravel; granite, gneissic granite and gneiss; and quartzite, slate, phyllite, greywacke, hornfels and schist. Sandy yellow earths.	Mainly granite, gneissic granite and gneiss. Also quartzite, slate, phyllite, greywacke, hornfels and schist. Sandy granite soils.		
LOCATION EXAMPLE	NW of Annandale South	SE of Annandale South	Woomargama State Forest		Mount Narra Narra
TREES > 8 m	<i>Acacia dealbata</i> - Silver Wattle <i>Eucalyptus blakelyi</i> - Blakely's Red Gum <i>E. bridgesiana</i> - Apple Box <i>E. melliodora</i> - Yellow Box	<i>Acacia dealbata</i> Silver Wattle <i>A. melanoxylon</i> Blackwood <i>Brachychiton populneus</i> Kurrajong <i>Eucalyptus albens</i> White Box <i>E. blakelyi</i> Blakely's Red Gum <i>E. bridgesiana</i> Apple Box <i>E. macrorhyncha</i> Red Stringybark <i>E. mannifera</i> Brittle Gum <i>E. melliodora</i> Yellow Box <i>E. polyanthemos</i> Red Box	<i>Acacia dealbata</i> <i>A. doratoxylon</i> <i>A. implexa</i> <i>Allocasuarina verticillata</i> <i>Brachychiton populneus</i> <i>Callitris endlicheri</i> <i>Eucalyptus blakelyi</i> <i>E. goniocalyx</i> <i>E. macrorhyncha</i> <i>E. nortonii</i> <i>E. polyanthemos</i> <i>Exocarpos cupressiformis</i>	Silver Wattle Currawang Hickory Wattle Drooping Sheoak Kurrajong Black Cypress Pine Blakely's Red Gum Long-leaf Box Red Stringybark Silver Bundy Red Box Native Cherry	<i>Acacia dealbata</i> <i>A. melanoxylon</i> <i>Eucalyptus bicostata</i> + <i>E. camphora</i> <i>E. dives</i> <i>E. goniocalyx</i> <i>E. macrorhyncha</i> <i>E. mannifera</i> <i>E. nortonii</i> <i>E. robertsonii</i> <i>E. rubida</i> <i>E. viminalis</i> <i>Exocarpos cupressiformis</i> + creeks/soaks
SHRUBS 1.5 - 8 m	# <i>Bursaria spinosa</i> - Native Blackthorn <i>Kunzea ericoides</i> - Burgan + <i>Leptospermum continentale</i> - Prickly Tea-tree # not noted in area but suggested for replanting + soaks/poorly drained sites	<i>Acacia paradoxa</i> - Kangaroo Thorn # <i>A. rubida</i> - Red-stemmed Wattle # <i>Bursaria spinosa spinosa</i> - Sweet Bursaria <i>Kunzea ericoides</i> - Burgan + <i>Leptospermum continentale</i> - Prickly Tea-tree <i>Pultenaea spinosa</i> - Grey Bush-pea + soaks/poorly drained sites	<i>Acacia gunnii</i> - Ploughshare Wattle <i>A. paradoxa</i> - Kangaroo Thorn <i>A. rubida</i> - Red-stemmed Wattle <i>A. verniciflua</i> - Varnish Wattle <i>Bursaria spinosa</i> - Native Blackthorn <i>Calytrix tetragona</i> - Common Fringe-myrtle <i>Cassinia aculeata</i> - Common Cassinia <i>Daviesia latifolia</i> - Hop Bitter-pea <i>D. leptophylla</i> - Slender Bitter-pea <i>Dodonaea viscosa angustissima</i> -	Narrow-leaf Hop-bush <i>I. australis</i> - Austral Indigo <i>Kunzea parvifolia</i> - Violet Kunzea + <i>Leptospermum continentale</i> - Prickly Tea-tree <i>Pultenaea spinosa</i> - Grey-Bush-pea <i>P. procumbens</i> - Heathy Bush-pea <i>Persoonia rigida</i> - Hairy Geebung + soaks/poorly drained sites	<i>Acacia rubida</i> <i>Bursaria spinosa</i> <i>Cassinia aculeata</i> <i>Kunzea ericoides</i> + <i>Leptospermum continentale</i> <i>Persoonia rigida</i> + soaks/poorly drained sites
GROUND COVERS	+ <i>Blechnum nudum</i> - Water Fern <i>Bothriochloa macra</i> - Red Grass + <i>Carex</i> spp. - Sedge <i>Juncus</i> spp. - Rush <i>Microlaena stipoides</i> - Weeping Grass + <i>Phragmites australis</i> - Common Reed	<i>Poa</i> spp. - Tussock Grasses <i>Rytidosperma</i> spp. - Wallaby Grass <i>Themeda triandra</i> - Kangaroo Grass + <i>Typha</i> spp. - Cumbungi + creeks/soaks/poorly drained sites	<i>Adiantum aethiopicum</i> - Common Maidenhair * <i>Brachyloma daphnoides</i> - Daphne Heath <i>Bulbine bulbosa</i> - Bulbine Lily * <i>Chrysocephalum apiculatum</i> - Yellow Buttons * <i>Dillwynia sericea</i> - Showy Parrot-pea <i>Glycine clandestina</i> - Twining Glycine * <i>Hardenbergia violacea</i> - Purple Coral Pea	<i>Hibbertia obtusifolia</i> - Grey Guinea-flower <i>Isotoma axillaris</i> - Showy Isotome * <i>Melichrus urceolatus</i> - Urn Heath <i>Microlaena stipoides</i> - Weeping Grass <i>Pimelea</i> spp. - Rice-flower <i>Poa sieberiana</i> - Fine-leaf Tussock Grass <i>Rytidosperma pallidum</i> - Red-anther Wallaby Grass	* <i>Rytidosperma</i> spp. - Wallaby Grass <i>Stypantra glauca</i> - Nodding Blue-lily <i>Themeda triandra</i> - Kangaroo Grass <i>Xanthorrhoea</i> spp. - Grass-tree * <i>Xerochrysum</i> spp. - Everlasting * mainly N and NW aspect